Laparoscopic Ventral Hernia Repair with Mesh (Laparoscopic repair of abdominal hernia with plastic)

Dr. Sitara Kommareddi
6567 East Carondelet Drive Suite 435
Tucson, Arizona 85710
520-512-5757 (office)

520-570-1278 (answering service)

Follow Up:

- Call in next several days to make an appointment to see me in 2-3 weeks.
- Call if you have Temp > 102, persistent, vomiting, worsening pain, redness that rapidly worsens or if you are unable to urinate 6-8 hours after you arrive home.

520-882-3211 (fax)

<u>Prescriptions</u>: Please give at least 2 days notice to obtain refills on meds after surgery. **Supplies to get:** a few small Band-Aids to use if any oozing from wounds, Ibuprofen 200mg tabs, Tylenol 500mg tabs, colace (a stool softener).

<u>Postoperative expectations/Restrictions:</u>

PAIN IS THE MAIN ISSUE AFTER THIS PROCEDURE. SOME PATIENTS HAVE SEVERE ACUTE PAIN OR RARELY LATER CHRONIC PAIN THAT MAY REQUIRE MORE AGGRESSIVE OPTIONS SUCH AS NERVE INJECTION/RESECTION OR LONG TERM PAIN MEDS. THIS IS AN INHERENT RISK OF THIS PROCEDURE.

THERE IS NO PERFECT HERNIA REPAIR. ALL APPROACHES HAVE A RISK OF RECURRENCE AND COMPLICATIONS. YOU MUST READ ONLINE COMMENTS ABOUT PATIENT EXPERIENCES.

- Wound pain is significant for the first 2 weeks to one month after this technique of repairing hernias. It will be present to varying degrees for several months and may require aggressive medication. You may also have some of the same symptoms you've had before surgery. These will take a few day/weeks/ months to recover. The pain may be burning, stabbing, aching, hot/cold or numb/tingling. As you return to normal activities over time, these symptoms may reoccur periodically. Many patients have occasional twinges for years.
- You will have bruising, swelling, hard areas in wounds and over the original hernia site that will resolve over several weeks to months
- **Diet**: As usual.
- Fatigue, nausea, bloating, constipation may occur.
- Alcohol should be avoided for the first 5 days.

- Activity: No lifting more than 15 lbs. for 4 weeks. Avoid swimming, running, golf, yoga, Pilates, sit-ups etc. for that period of time. After 4 weeks, you may slowly resume to your pace as your symptoms allow.
- **Urinary retention**: Rarely patients are unable to urinate. If you are unable to urinate 6 hours after you get home, please call. I may have you present to an E.R. to have a bladder check, and possibly place a 12 French Bladder catheter to stay in place for 5 days and remove in my clinic.
- Travel should be avoided for 2 weeks-4 weeks after the procedure, if possible.
- **Driving** should be avoided for 7-14days. Subsequently drive if you are fully alert, not on narcotics, and feel you can appropriately react to any driving stressor.
- **Home medications** may be resumed as usual upon arrival home. Specific instructions may be given regarding certain medications such as blood thinners.

Wound Care:

- You will have large white pads over wound. Remove the morning after surgery, and discard.
- White strips on wounds (steristrips) will fall off, or you may remove 2 weeks after surgery or I will remove.
- You may have bloody, pink, yellow, orange discharge from the wounds. In that case place band aids as needed.
- Shower: Shower the morning after procedure. Do not soak in water for 3 weeks.

For Pain relief: Take meds with food/water

- Abdominal binder, wear as much as it helps for 2 weeks then as needed. You can put in washer/dryer to soften.
- Ice on the wounds 20 minutes at a time, 2-3 x a day is helpful
- **#1.** Tylenol (acetaminophen) 1000mg every 8 hours by mouth. **Try to take it regularly for 3 days, then only as needed.** (Avoid if cirrhosis). Obtain at drugstore, no need for prescription.

#2 Ibuprofen 600 mg every 8 hours by mouth. **Try to take it regularly for 3 days, then only as needed**. (Avoid if gastritis, ulcers, inflammatory bowel disease, intolerance to this drug, kidney failure/kidney removal, or if on blood thinners). Obtain at drugstore, no need for a prescription.

ALTERNATE IBUPROFEN AND TYLENOL EVERY 4 HOURS.

#3. Pregabalin (Lyrica) to take by mouth every night until all tabs gone. This is a nerve med to treat pain. It may cost around \$30.00. If your insurance does not cover, please consider paying out of pocket. You will receive a prescription.

#4 A narcotic prescription may be provided. You can take in addition to all of those above.

For Nausea relief:

#1 Ondansetron 4 mg every 4 hours as needed for nausea. You will receive a prescription.

#2 Benadryl 25 mg tablet by mouth every 8 hours if #1 not enough. This may be obtained at the drugstore without prescription.

Treatment of Constipation:

If you do not have a bowel movement within 24 hours after surgery, then:

#1 Colace 100 mg by mouth in am and pm. If not enough to help in 24 hours, then stay on it and add:

#2 Milk of Magnesia 30 ml once a day. If no B.M. 24 hours later, then stop it, stay on Colace and add:

#3 Magnesium Citrate 1 bottle once by mouth. Take ice cold over 1 hour, poured over ice along with large glass of cold water. If no B.M. in 24 hours:

#4 Repeat magnesium citrate 1 bottle by mouth once again and take a Fleets enema.

#5 If none of the above work call me for other options.

All may be obtained at the drugstore without a prescription.

Itching:

#1 Benadryl 25 mg tablet by mouth every 8 hours.

#2 Once steristrips off of wounds, try Benadryl ointment or topical steroid ointment to areas of itching 3-4 x a day. Can get both at drugstore without prescription.

Blistering at wound site:

White strips on wounds (steristrips) can cause blistering or redness in some patients. . If this occurs you may remove them and let me know. If not helpful, please call.

Time off work/ school:

- **For desk jobs**: Plan on taking at least 7-14 days off. You may return once you feel you can work a full day as long as you follow restrictions. **We will discuss specifics for other employment.**
- Please let me know if you need a work release. You may get it on day of follow up or we can fax
 to number you provide. Please be specific as to when you want to go back to work and if any
 restrictions.
- Any disability paperwork may be dropped off. We may take 5-7 days to complete. There is a fee of \$25.00.

C 07/R22